

# Wildlife Soybean

## Eagle Forage Soybean RR™ Annual

**Large Lad RR™** soybeans are our most popular food plot variety and are known for the excellent seed yield and tonnage production. Large Lad plants can reach heights up to 84 inches and provide excellent forage for deer and cattle. Easy to grow, with resistance to many foliar diseases, root rots, stem canker and races of nematodes, they are widely adapted to numerous soil types, and will withstand harsher growing conditions than regular soybeans. Planting Large Lad RR™ will supply more green food all season for your deer and bird hunting.

**Wildlife Manager's Mix RR™** soybeans originally developed for our northern customers, Manager's Mix RR is a blend containing both forage types, and climbing soybeans. Featuring 4 maturity groups which mature at different intervals and allow the deer to feed on both mature soybeans and green leaf tissue. High tonnage, excellent browse tolerance and are relatively easy to grow, resistance to many foliar diseases, root rots, stem canker and races of nematodes, and will withstand harsher growing conditions than regular soybeans. Planting Wildlife Manager's Mix RR™ will supply more green food all season for your deer and bird hunting.



**Soil Types:** Wide variety of soils optimum pH is from 6.0 to 7.0.

**Maintenance:** Low maintenance

**Seeding Rate:** 50-80 lbs. per acre

**Frost Seeding:** Not applicable

**Planting Date:** April - June

**Planting Depth:** ¾ - 1 inch





8496 Swan Pond Rd \* PO Box 308  
Weiner, AR 72479 \* 870-684-7377

## Glyphosate Tolerant Soybean Planting Tips

Row Spacing	140,000	175,000	200,000
	Seeds per row foot for each population desired		
38" row	10.2	12.7	14.5
36" row	9.6	12.1	13.8
30" row	8.0	10.0	11.5
20" row	5.4	6.7	7.7
15" row	4.0	5	5.7*
7.5" row	2.0	2.5	2.9*
Broadcast/ square foot	3.2	4.0	4.6*
RECOM- MENDED SOIL CONDITIONS AND DEER DENSITY	Use this rate for good soil condi- tions or low deer density	Use this rate for normal deer density	*Use this rate or higher for less than ideal soil and/or heavy deer density

\*Data compiled from multiple states and sources including extension and university data.

©2016. Eagle Seed Company. All rights reserved.

### FERTILIZER:

- The optimum pH for soybeans is between 6.0 and 7.0. 6.5 is ideal.
- Inoculate soybeans with *Bradyrhizobium japonicum* if the soil has either high or low pH, has been water-logged or extremely cold, or has no recent soybean production. Inoculant can also reduce your costs. Nitrogen is not needed on decent soil if the soybeans are inoculated. Starter N is fine. We recommend inoculant be applied to the seed just prior to planting. Add the inoculant to the seed hopper and stir gently, do not add moisture (check with inoculant label).
- For best results, fertilize according to your farm's soil test as analyzed by your State Extension Service. If you don't have a soil test, use inoculant and 0-40-70 (units). Soybeans remove a minimum of 40 lbs of Phosphorous and 70 lbs of Potassium. For maximum deer antler and body weight production, Dr. G. Woods, deer biologist, uses 0-70-120 (units).
- Always store seed in a low humidity environment below 90°F.

### INTER-SEEDING WITH ROUNDUP READY CORN:

To maximize plant growth for both species, it is best to plant soybean and corn in separate patches. If you prefer inter-seeding them, studies show that optimum results are achieved with 3-5 lbs. corn and one bag per acre of soybeans. Add N for the corn, soybeans will fix only enough for their own use.

### HERBICIDE APPLICATION:

- It is best to start with a clean, weed-free seed bed. Use tillage or a pre-plant burn-down herbicide such as Roundup (glyphosate only version), Liberty, or Gramoxone depending on your emerged weeds. Pre-emerge herbicides such as Valor, Metribuzin, Treflan or Prowl can be used at their labeled rate. Add Fomesafen for increased pigweed control over the top at the labeled rate.
- Roundup (Glyphosate-only Roundup or generic glyphosate labeled for soybean) can be applied to the crop at any time, but weeds need to be controlled in the first 3 to 4 weeks to maximize soybean plant growth. It is important to spray weeds BEFORE they reach 4 inches in height.
- Roundup PowerMax and WeatherMax are the best products; the only active ingredient should be glyphosate, though most formulations will also contain a surfactant (make sure to read and follow label directions).

### PLANTING TIME, DEPTH, SPACINGS:

- In general, most northern food plotters can plant in late May or early June. Southern food plots can usually be planted in late April to early May. Check with your local extension service for optimum dates. This is usually after your last spring frost when the soil temperature at a one inch planting depth is above 65°F, but no greater than 90°F. Soybeans have a different kind of emergence, so it is usually best to plant after a rain. Rain can cause surface soil compaction and crusting which prevents plant establishment.
- The recommended planting depth is 3/4 to 1 inch into a moist soil. Do not plant in a dry soil, or in muddy conditions. Put seeds 1/4 inch below soil moisture line, but no deeper than 1.5 inches. If you do not see emergence in 7 days, check the seeds for sprouting. When broadcasting, increase your seeding rate to 75 lbs per acre since not every seed will have proper seed to soil contact.

Seed size will vary, so use a minimum of one bag to the acre. The chart is provided to calibrate your drill or planter based on row width. If planting in less than optimal conditions or heavy deer densities, use the higher plant population and a narrower row, up to 15 inches in width.

### ESTABLISHMENT:

Plants grow slower in cooler spring temperatures, so use deer deterrents for the first 4-8 weeks, especially in areas of heavy deer pressure or in cooler temperatures.

### DRAINAGE:

Make sure your crops have good drainage. Soybean plant growth and nitrogen fixation can be lost when water stands on an area for 3 or more days. Avoid flood prone areas.

# Wildlife Manager's Mix RR™ Soybeans

Wildlife Food Plot Soybean

